

GLOBAL EXISTENCE OF SMOOTH SOLUTIONS OF A 3D LOGLOG ENERGY-SUPERCritical WAVE EQUATION

TRISTAN ROY

ABSTRACT. We prove global existence of smooth solutions of the 3D loglog energy-supercritical wave equation $\partial_{tt}u - \Delta u = -u^5 \log^c(\log(10 + u^2))$ with $0 < c < \frac{8}{225}$ and smooth initial data $(u(0) = u_0, \partial_t u(0) = u_1)$. First we control the $L_t^4 L_x^{12}$ norm of the solution on an arbitrary size time interval by an expression depending on the energy and an *a priori* upper bound of its $L_t^\infty \tilde{H}^2(\mathbb{R}^3)$ norm, with $\tilde{H}^2(\mathbb{R}^3) := \dot{H}^2(\mathbb{R}^3) \cap \dot{H}^1(\mathbb{R}^3)$. The proof of this long time estimate relies upon the use of some potential decay estimates [1, 12] and a modification of an argument in [17]. Then we find an *a posteriori* upper bound of the $L_t^\infty \tilde{H}^2(\mathbb{R}^3)$ norm of the solution by combining the long time estimate with an induction on time of the Strichartz estimates.

1. INTRODUCTION

We shall consider the defocusing loglog energy-supercritical wave equation

$$(1.1) \quad \partial_{tt}u - \Delta u = -f(u)$$

where $u : \mathbb{R} \times \mathbb{R}^3 \rightarrow \mathbb{R}$ is a real-valued scalar field and $f(u) := u^5 g(u)$ with $g(u) := \log^c(\log(10 + u^2))$, $0 < c < \frac{8}{225}$. *Classical solutions* of (1.1) are solutions that are infinitely differentiable and compactly supported in space for each fixed time t . It is not difficult to see that *classical solutions* of (1.1) satisfy the energy conservation law

$$(1.2) \quad E := \frac{1}{2} \int_{\mathbb{R}^3} (\partial_t u(t, x))^2 dx + \frac{1}{2} \int_{\mathbb{R}^3} |\nabla u(t, x)|^2 dx + \int_{\mathbb{R}^3} F(u(t, x)) dx$$

where $F(u) := \int_0^u f(v) dv$. *Classical solutions* of (1.1) enjoy three symmetry properties that we use throughout this paper

- the *time translation invariance*: if u is a solution of (1.1) and t_0 is a fixed time then $\tilde{u}(t, x) := u(t - t_0, x)$ is also a solution of (1.1).
- the *space translation invariance*: if u is a solution of (1.1) and x_0 is a fixed point lying in $\mathbb{R}^3$ then $\tilde{u}(t, x) := u(t, x - x_0)$ is also a solution of (1.1)
- the *time reversal invariance*: if u is a solution to (1.1) then $\tilde{u}(t, x) := u(-t, x)$ is also a solution to (1.1)

The defocusing loglog energy-supercritical wave equation (1.1) is closely related to the power-type defocusing wave equations, namely

$$(1.3) \quad \partial_{tt}u - \Delta u = -|u|^{p-1}u$$

Solutions of (1.3) have an invariant scaling

$$(1.4) \quad u(t, x) \rightarrow u^\lambda(t, x) := \frac{1}{\lambda^{\frac{2}{p-1}}} u\left(\frac{t}{\lambda}, \frac{x}{\lambda}\right)$$

and (1.3) is $s_c := \frac{3}{2} - \frac{2}{p-1}$ critical. This means that the $\dot{H}^{s_c}(\mathbb{R}^3) \times \dot{H}^{s_c-1}(\mathbb{R}^3)$ norm of $(u(0), \partial_t u(0))$ is invariant under scaling or, in other words, $\|u^\lambda(0)\|_{\dot{H}^{s_c}(\mathbb{R}^3)} = \|u(0)\|_{\dot{H}^{s_c}(\mathbb{R}^3)}$ and $\|\partial_t u^\lambda(0)\|_{\dot{H}^{s_c-1}(\mathbb{R}^3)} = \|\partial_t u(0)\|_{\dot{H}^{s_c-1}(\mathbb{R}^3)}$. If $p = 5$ then $s_c = 1$ and this is why the quintic defocusing cubic wave equation

$$(1.5) \quad \partial_{tt} u - \Delta u = -u^5$$

is called the energy-critical equation. If $1 < p < 5$ then $s_c < 1$ and (1.3) is energy-subcritical while if $p > 5$ then $s_c > 1$ and (1.3) is energy-supercritical. Notice that for every $p > 5$ there exists two positive constant $\lambda_1(p)$, $\lambda_2(p)$ such that

$$(1.6) \quad \lambda_1(p)|u|^5 \leq |f(u)| \leq \lambda_2(p) \max(1, |u|^p)$$

This is why (1.1) is said to belong to the group of barely supercritical equations. There is another way to see that. Notice that a simple integration by part shows that

$$(1.7) \quad F(u) \sim \frac{u^6}{6} g(u)$$

and consequently the nonlinear potential term of the energy $\int_{\mathbb{R}^3} F(u) dx \sim \int_{\mathbb{R}^3} u^6 g(u) dx$ just barely fails to be controlled by the linear component, in contrast to (1.5).

The energy-critical wave equation (1.5) has received a great deal of attention. Grillakis [4, 5] established global existence of smooth solutions (global regularity) of this equation with smooth initial data $(u(0) = u_0, \partial_t u(0) = u_1)$. His work followed the work of Rauch for small data [10] and the one of Struwe [15] handling the spherically symmetric case. Later Shatah and Struwe [12] gave a simplified proof of this result. Kapitanski [6] and, independently, Shatah and Struwe [13], proved global existence of solutions with data (u_0, u_1) in the energy class.

We are interested in proving global regularity of (1.1) with smooth initial data (u_0, u_1) . By standard persistence of regularity results it suffices to prove global existence of solutions $u \in \mathcal{C}\left([0, T], \tilde{H}^2(\mathbb{R}^3)\right) \cap \mathcal{C}^1\left([0, T], H^1(\mathbb{R}^3)\right)$ with data $(u_0, u_1) \in \tilde{H}^2(\mathbb{R}^3) \times H^1(\mathbb{R}^3)$. Here $\tilde{H}^2(\mathbb{R}^3)$ denotes the following space

$$(1.8) \quad \tilde{H}^2(\mathbb{R}^3) := \dot{H}^2(\mathbb{R}^3) \cap \dot{H}^1(\mathbb{R}^3)$$

In view of the local well-posedness theory [8], standard limit arguments and the finite speed of propagation it suffices to find an *a priori* upper bound of the form

$$(1.9) \quad \|(u(T), \partial_t u(T))\|_{\tilde{H}^2(\mathbb{R}^3) \times H^1(\mathbb{R}^3)} \leq C_1 \left(\|u_0\|_{\tilde{H}^2(\mathbb{R}^3)}, \|u_1\|_{H^1(\mathbb{R}^3)}, T \right)$$

for all times $T > 0$ and for *classical solutions* u of (1.1) with smooth and compactly supported data (u_0, u_1) . Here C_1 is a constant depending only on $\|u_0\|_{\tilde{H}^2(\mathbb{R}^3)}$, $\|u_1\|_{H^1(\mathbb{R}^3)}$ and the time T .

The global behaviour of the solutions of the supercritical wave equations is poorly understood, mostly because of the lack of conservation laws in $\tilde{H}^2(\mathbb{R}^3)$. Nevertheless

Tao [17] was able to prove global regularity for another barely supercritical equation, namely

$$(1.10) \quad \partial_{tt}u - \Delta u = -u^5 \log(2 + u^2)$$

with radial data. The main result of this paper is

Theorem 1.1. *The solution of (1.1) with smooth data (u_0, u_1) exists for all time. Moreover there exists a nonnegative constant $M_0 = M_0(\|u_0\|_{\tilde{H}^2(\mathbb{R}^3)}, \|u_1\|_{H^1(\mathbb{R}^3)})$ depending only on $\|u_0\|_{\tilde{H}^2(\mathbb{R}^3)}$ and $\|u_1\|_{H^1(\mathbb{R}^3)}$ such that*

$$(1.11) \quad \|u\|_{L_t^\infty \tilde{H}^2(\mathbb{R} \times \mathbb{R}^3)} + \|\partial_t u\|_{L_t^\infty H^1(\mathbb{R} \times \mathbb{R}^3)} \leq M_0$$

We recall some basic properties and estimates. Let Q be a function, let J be an interval and let $t_0 \in J$ be a fixed time. If u is a *classical solution* of the more general problem $\partial_{tt}u - \Delta u = Q$ then u satisfies the Duhamel formula

$$(1.12) \quad u(t) = u_{l,t_0}(t) + u_{nl,t_0}(t), \quad t \in J$$

with u_{l,t_0} , u_{nl,t_0} denoting the linear part and the nonlinear part respectively of the solution starting from t_0 . Recall that

$$(1.13) \quad u_{l,t_0}(t) = \cos(t - t_0)Du(t_0) + \frac{\sin(t - t_0)D}{D}\partial_t u(t_0)$$

and

$$(1.14) \quad u_{nl,t_0}(t) = - \int_{t_0}^t \frac{\sin(t - t')D}{D} Q(t') dt'$$

with D the multiplier defined by $\widehat{Df}(\xi) := |\xi| \widehat{f}(\xi)$. An explicit formula for $\frac{\sin(t - t')D}{D} Q(t')$ and $t \neq t'$ is

$$(1.15) \quad \left[\frac{\sin(t - t')D}{D} Q(t') \right] (x) = \frac{1}{4\pi|t - t'|} \int_{|x - x'| = |t - t'|} Q(t', x') dS(x')$$

For a proof see for example [14]. We recall that u_{l,t_0} satisfies $\partial_{tt}u_{l,t_0} - \Delta u_{l,t_0} = 0$, $u_{l,t_0}(t_0) = u(t_0)$ and $\partial_t u_{l,t_0}(t_0) = \partial_t u(t_0)$ while u_{nl,t_0} is the solution of $\partial_{tt}u_{nl,t_0} - \Delta u_{nl,t_0} = Q$, $u_{nl,t_0}(t_0) = 0$ and $\partial_t u_{nl,t_0}(t_0) = 0$. We recall the Strichartz estimate [3, 7, 8, 14]

$$(1.16) \quad \|u\|_{L_t^q L_x^r(J \times \mathbb{R}^3)} \lesssim \|\partial_t u(t_0)\|_{L_x^2(\mathbb{R}^3)} + \|\nabla u(t_0)\|_{L_x^2(\mathbb{R}^3)} + \|Q\|_{L_t^1 L_x^2(J \times \mathbb{R}^3)}$$

if (q, r) is wave admissible, i.e $(q, r) \in (2, \infty] \times [2, \infty]$ and $\frac{1}{q} + \frac{3}{r} = \frac{1}{2}$.

We set some notation that appear throughout the paper. We write $C = C(a_1, \dots, a_n)$ if C only depends on the parameters $a_1, \dots, a_n$. We write $A \lesssim B$ if there exists a universal nonnegative constant $C' > 0$ such that $A \leq C' B$. $A = O(B)$ means $A \lesssim B$. More generally we write $A \lesssim_{a_1, \dots, a_n} B$ if there exists a nonnegative constant $C' = C(a_1, \dots, a_n)$ such that $A \leq C' B$. We say that C'' is the constant determined by $\lesssim$ in $A \lesssim_{a_1, \dots, a_n} B$ if C'' is the smallest constant among the C' s

such that $A \leq C' B$. We write $A \ll_{a_1, \dots, a_n} B$ if there exists a universal nonnegative small constant $c = c(a_1, \dots, a_n)$ such that $A \leq cB$. Similar notions are defined for $A \gtrsim B$, $A \gtrsim_{a_1, \dots, a_n} B$ and $A \gg B$. In particular we say that C'' is the constant determined by $\gtrsim$ in $A \gtrsim B$ if C'' is the largest constant among the C' 's such that $A \geq C' B$. If x is number then $x+$ and $x-$ are slight variations of x : $x+ := x + \alpha\epsilon$ and $x- := x - \beta\epsilon$ for some $\alpha > 0$, $\beta > 0$ and $0 < \epsilon \ll 1$.

Let Γ_+ denote the forward light cone

$$(1.17) \quad \Gamma_+ = \{(t, x) : t > |x|\}$$

and if $J = [a, b]$ is an interval let $\Gamma_+(J)$ denote the light cone truncated to J i.e

$$(1.18) \quad \Gamma_+(J) := \Gamma_+ \cap (J \times \mathbb{R}^3)$$

Let $e(t)$ denote the local energy i.e

$$(1.19) \quad e(t) := \frac{1}{2} \int_{|x| \leq t} (\partial_t u(t, x))^2 dx + \frac{1}{2} \int_{|x| \leq t} |\nabla u(t, x)|^2 dx + \int_{|x| \leq t} F(u(t, x)) dx$$

If u is a solution of (1.1) then by using the finite speed of propagation and the Strichartz estimates we have

$$(1.20) \quad \|u\|_{L_t^q L_x^r(\Gamma_+(J))} \lesssim \|\nabla u(b)\|_{L_x^2(\mathbb{R}^3)} + \|\partial_t u(b)\|_{L_x^2(\mathbb{R}^3)} + \|Q\|_{L_t^1 L_x^2(\Gamma_+(J))}$$

if (q, r) is wave admissible. If $J_1 := [a_1, a_2]$ and $J_2 := [a_2, a_3]$ then we also have

$$(1.21) \quad \|u\|_{L_t^q L_x^r(\Gamma_+(J_1))} \lesssim \|\nabla u(a_3)\|_{L_x^2(\mathbb{R}^3)} + \|\partial_t u(a_3)\|_{L_x^2(\mathbb{R}^3)} + \|Q\|_{L_t^1 L_x^2(\Gamma_+(J_1 \cup J_2))}$$

We recall also the well-known Sobolev embeddings. If h is a smooth function then

$$(1.22) \quad \|h\|_{L^\infty(\mathbb{R}^3)} \lesssim \|h\|_{\dot{H}^2(\mathbb{R}^3)}$$

and

$$(1.23) \quad \|h\|_{L^6(\mathbb{R}^3)} \lesssim \|\nabla h\|_{L^2(\mathbb{R}^3)}$$

If u is the solution of (1.1) with data $(u_0, u_1) \in \tilde{H}^2(\mathbb{R}^3) \times H^1(\mathbb{R}^3)$ then we get from (1.22)

$$(1.24) \quad E \lesssim \|u_0\|_{\tilde{H}^2(\mathbb{R}^3)}^2 \max\left(1, \|u_0\|_{\tilde{H}^2(\mathbb{R}^3)}^4 g(\|u_0\|_{\tilde{H}^2(\mathbb{R}^3)})\right)$$

We shall use the Paley-Littlewood technology. Let $\phi(\xi)$ be a bump function adapted to $\{\xi \in \mathbb{R}^3 : |\xi| \leq 2\}$ and equal to one on $\{\xi \in \mathbb{R}^3 : |\xi| \leq 1\}$. If $(M, N) \in 2^{\mathbb{Z}} \times 2^{\mathbb{Z}}$ are dyadic numbers then the Paley-Littlewood projection operators P_M , $P_{<N}$ and $P_{\geq N}$ are defined in the Fourier domain by

$$(1.25) \quad \widehat{P_M f}(\xi) := \left(\phi\left(\frac{\xi}{M}\right) - \phi\left(\frac{\xi}{2M}\right)\right) \hat{f}(\xi)$$

$$(1.26) \quad \widehat{P_{<N}f}(\xi) := \sum_{M < N} \widehat{P_M f}(\xi)$$

and

$$(1.27) \quad \widehat{P_{\geq N}f}(\xi) := \sum_{M \geq N} \widehat{P_M f}(\xi)$$

The inverse Sobolev inequality can be stated as follows

Proposition 1. "Inverse Sobolev inequality" [16] *Let g be a smooth function such that $\|g\|_{\dot{H}^1(\mathbb{R}^3)} \lesssim E^{\frac{1}{2}}$ and $\|P_{\geq N}g\|_{L_x^6(\mathbb{R}^3)} \gtrsim \eta$ for some real number $\eta > 0$ and for some dyadic number $N > 0$. Then there exists a ball $B(x, r) \subset \mathbb{R}^3$ with $r = O(\frac{1}{N})$ such that we have the mass concentration estimate*

$$(1.28) \quad \int_{B(x, r)} |g(y)|^2 dy \gtrsim \eta^3 E^{-\frac{1}{2}} r^2$$

We also recall a result that shows that the mass of solutions of (1.1) can be locally in time controlled

Proposition 2. "Local mass is locally stable" [16] *Let J be a time interval, let $t, t' \in J$ and let $B(x, r)$ be a ball. Let u be a solution of (1.1). Then*

$$(1.29) \quad \left(\int_{B(x, r)} |u(t', y)|^2 dy \right)^{\frac{1}{2}} = \left(\int_{B(x, r)} |u(t, y)|^2 dy \right)^{\frac{1}{2}} + O\left(E^{\frac{1}{2}} |t - t'|\right)$$

Notice that this result, proved for (1.5) in [16], is also true for (1.1). Indeed the proof relied upon the fact that the $L^2(\mathbb{R}^3)$ norm of the velocity of the solution of (1.5) at time t is bounded by the square root of its energy, which is also true for the solution of (1.1) (by (1.2) and (1.7)).

Now we make some comments with respect to Theorem 1.1. If the function g were a positive constant, then it would be easy to prove that the solution of (1.1) with data $(u_0, u_1) \in \tilde{H}^2(\mathbb{R}^3) \times H^1(\mathbb{R}^3)$, since we have a good global theory for (1.5). Therefore we can hope to prove global well-posedness for g slowly increasing to infinity, by extending the technology to prove global well-posedness for (1.5). Notice also that Tao, in [16], found that the solution u of (1.5) satisfies

$$(1.30) \quad \|u\|_{L_t^4 L_x^{12}(\mathbb{R} \times \mathbb{R}^3)} \lesssim \tilde{E}^{\tilde{E}^{O(1)}}$$

with $\tilde{E}$ the energy of u . The structure of g is a double log: it is, roughly speaking, the inverse function of the tower exponential bound in (1.30).

Now we explain the main ideas of this paper.

In [16], Tao was able to bound on arbitrary long time intervals the $L_t^4 L_x^{12}$ norm of solutions of the energy-critical equation (1.5) by a quantity that depends exponentially on their energy. This estimate can be viewed as a long time estimate. Unfortunately we cannot expect to prove a similar result for (1.1) since we are not in the energy-critical regime. However we shall prove the following proposition

Proposition 3. "Long time estimate"

Let $J = [t_1, t_2]$ be a time interval. Let u be a classical solution of (1.1). Assume that

$$(1.31) \quad \|u\|_{L_t^\infty \tilde{H}^2(J \times \mathbb{R}^3)} \leq M$$

holds for some $M \geq 0$. Then there exist three constants $C_{L,0} > 0$, $C_{L,1} > 0$ and $C_{L,2} > 0$ such that

- if $E < \frac{1}{g^{\frac{1}{2}}(M)}$ (small energy regime) then

$$(1.32) \quad \|u\|_{L_t^4 L_x^{12}(J \times \mathbb{R}^3)}^4 \leq C_{L,0}$$

- if $E \gtrsim \frac{1}{g^{\frac{1}{2}}(M)}$ (large energy regime) then

$$(1.33) \quad \|u\|_{L_t^4 L_x^{12}(J \times \mathbb{R}^3)}^4 \leq (C_{L,1}(Eg(M)))^{C_{L,2}(E^{\frac{193}{4}} + g^{\frac{225}{8}}(M))}$$

This proposition shows that we can control the $L_t^4 L_x^{12}(J \times \mathbb{R}^3)$ norm of solutions of (1.1) by their energy and an *a priori* bound of their $L_t^\infty \tilde{H}^2(J \times \mathbb{R}^3)$ norm. We would like to control the pointwise-in-time $\tilde{H}^2(\mathbb{R}^3) \times H^1(\mathbb{R}^3)$ norm of u on an interval $[0, T]$, with T arbitrary large. This is done by an induction on time. We assume that this norm is controlled on $[0, T]$ by a number M_0 . Then by continuity we can find a slightly larger interval $[0, T']$ such that this norm is bounded by (say) $2M_0$ on $[0, T']$. This is our *a priori* bound. We subdivide $[0, T']$ into subintervals where the $L_t^4 L_x^{12}$ norm of u is small and we control the pointwise-in-time $\tilde{H}^2(\mathbb{R}^3) \times H^1(\mathbb{R}^3)$ norm of u on each of these subintervals (see Lemma 4). Since g varies slowly we can estimate the number of intervals of this partition by using Proposition 3 and we can prove *a posteriori* that $\|u(t)\|_{\tilde{H}^2(\mathbb{R}^3)} + \|\partial_t u(t)\|_{\tilde{H}^1(\mathbb{R}^3)}$ is bounded on $[0, T']$ by M_0 , provided that M_0 is large enough: see Section 2.

The proof of Proposition 3 is a modification of the argument used in [16] to establish a tower-exponential bound of the $L_t^4 L_x^{12}(J \times \mathbb{R}^3)$ norm of v , solution of (1.5). We divide J into subintervals J_i where the $L_t^4 L_x^{12}$ norm of u , solution of (1.1), is "substantial". Then by using the Strichartz estimates and the Sobolev embedding (1.22) we notice that the $L_t^\infty L_x^6(J_i \times \mathbb{R}^3)$ norm of u is also "substantial", more precisely we find a lower bound that depends on the energy E and $g(M)$. Then by Proposition 1 we can localize a bubble where the mass concentrates and we prove that the size of these subintervals is also "substantially" large. Tao [16] used the mass concentration to construct a solution $\tilde{v}$ of (1.5) that has a smaller energy than v and that coincides with v outside a cone. The idea behind that is to use an induction on the levels of energy, due to Bourgain [2], and the small energy theory following from the Strichartz estimates in order to control the $L_t^4 L_x^{12}$ norm of v outside a cone. Unfortunately it seems almost impossible to apply this procedure to our problem. Indeed the energy of the constructed solution $\tilde{u}$ is smaller than the energy E of u by an amount that depends on E but also on $g(M)$ and therefore an induction on the levels of the energy is possible if the $L_t^\infty \tilde{H}^2(J \times \mathbb{R}^3)$ norm of $\tilde{u}$ can be controlled by M , which is far from being trivial. It turns out that we do not need to use the Bourgain induction method. Indeed since we know that the size of the subintervals J_i is substantially large and since we have a good control of the $L_t^4 L_x^{12}$ norm on these subintervals it suffices to find an upper bound of the size of

their union in order to conclude. To this end we divide a cone containing the ball where the mass concentrates and the J_i s into truncated-in-time cones where the $L_t^4 L_x^{12}$ norm of u is “substantial”. Let $\tilde{J}_1, \tilde{J}_2, \dots$ be the sequence of time intervals resulting from this partition. The mass concentration helps us to control the size of the first time interval $\tilde{J}_1$. By using an asymptotic stability result we can prove, roughly speaking, that if we consider two successive subintervals $\tilde{J}_j, \tilde{J}_{j+1}$ resulting from this partition of the cone then the size of $\tilde{J}_{j+1}$ can be controlled by the size of $\tilde{J}_j$: see (3.35). But a potential energy decay estimate shows that if the size of the union of the J_i s is too large then we can find a large subinterval $[t'_1, t'_2]$ such that the $L_t^4 L_x^{12}$ norm of u on the cone truncated to $[t'_1, t'_2]$ is small. Therefore $[t'_1, t'_2]$ cannot be covered by many $\tilde{J}_j$ s and one of them is very large in comparison with its predecessor, which contradicts (3.35). At the end of the process we can find an upper bound of the size of the union of the subintervals J_i s and consequently we can control the $L_t^4 L_x^{12}$ norm of u on the interval J .

Remark 1.2. *Throughout the paper we frequently use the $x+$ and $x-$ notations. Indeed the point $(2, \infty)$ is not wave admissible. Therefore we will work with the point $(2+, \infty-)$: see (5.7) and (7.9). This generates slight variations of many quantities throughout this paper. Sometimes we might deal with quantities like $z := \frac{x+}{y-}$. We cannot conclude directly that $z = \frac{x}{y}+$. In this case we create a variation of y so small (comparing to that of x) that we have $z = \frac{x}{y}+$. These details have been omitted for the sake of readability. We strongly recommend that the reader ignores these slight variations at the first reading.*

Acknowledgements : The author would like to thank Terence Tao for suggesting him this problem.

2. PROOF OF THEOREM 1.1

The proof relies upon Proposition 3 and the following lemma that we prove in the next subsection.

Lemma 4. "Local boundedness" *Let $J = [t_1, t_2]$ be an interval. Assume that u is a classical solution of (1.1). Let $Z(t) := \|(u(t), \partial_t u(t))\|_{\tilde{H}^2(\mathbb{R}^3) \times H^1(\mathbb{R}^3)}$. There exists $0 < \epsilon < 1$ constant such that if*

$$(2.1) \quad \|u\|_{L_t^4 L_x^{12}(J \times \mathbb{R}^3)} \leq \frac{\epsilon}{g^{\frac{1}{4}}(Z(t_1))}$$

then there exists $C_l > 0$ such that

$$(2.2) \quad Z(t) \leq 2C_l Z(t_1)$$

for $t \in J$.

We claim that the following set

$$(2.3) \quad \mathcal{F} := \left\{ T \in [0, \infty) : \sup_{t \in [0, T]} \|(u(t), \partial_t u(t))\|_{\tilde{H}^2(\mathbb{R}^3) \times H^1(\mathbb{R}^3)} \leq M_0 \right\}$$

is equal to $[0, \infty)$ for some constant $M_0 := M_0(\|u_0\|_{\tilde{H}^2(\mathbb{R}^3)}, \|u_1\|_{H^1(\mathbb{R}^3)})$ large enough. Indeed

- $0 \in \mathcal{F}$: clear

- $\mathcal{F}$ is closed by continuity
- $\mathcal{F}$ is open. Indeed let $T \in \mathcal{F}$. Then by continuity there exists $\delta > 0$ such that

$$(2.4) \quad \sup_{t \in [0, T']} \| (u(t), \partial_t u(t)) \|_{H^2(\mathbb{R}^3) \times H^1(\mathbb{R}^3)} \leq 2M_0$$

for every $T' \in [0, T + \delta)$. By (1.32) and (1.33) we have

$$(2.5) \quad \|u\|_{L_t^4 L_x^{12}([0, T'] \times \mathbb{R}^3)}^4 \leq \max \left(C_{L,0}, (C_{L,1} E g(2M_0))^{C_{L,2} (E^{\frac{193}{4} + g} \frac{225}{8} + (2M_0))} \right)$$

Let $N \geq 1$ and let $\underline{Z}(0) := \max(Z(0), 1)$. Without loss of generality we can assume that $C_l \gg 1$ so that $2C_l \underline{Z}(0) \gg 1$ and $\log^c(2C_l \underline{Z}(0)) \gg 1$. We have, by the elementary rules of the logarithm¹ and the inequality $\log^c(2nx) \leq \log^c((2n)^x)$ for $n \geq 1$ and $x \gg 1$

$$(2.6) \quad \begin{aligned} \sum_{n=1}^N \frac{\epsilon^4}{g((2C_l)^n Z_0)} &\geq \sum_{n=1}^N \frac{\epsilon^4}{\log^c(\log((2C_l)^{2n} \underline{Z}^{2n}(0) + 10))} \\ &\gtrsim \sum_{n=1}^N \frac{1}{\log^c(2n \log(2C_l \underline{Z}(0)))} \\ &\gtrsim \frac{1}{\log^c(2C_l \underline{Z}(0))} \sum_{n=1}^N \frac{1}{\log^c(2n)} \\ &\gtrsim \frac{1}{\log^c(2C_l \underline{Z}(0))} \int_1^{N+1} \frac{1}{\log^c(2t)} dt \\ &\gtrsim \frac{1}{\log^c(2C_l \underline{Z}(0))} \int_1^{N+1} \frac{1}{t^{\frac{1}{2}}} dt \\ &\gtrsim \frac{N^{\frac{1}{2}}}{\log^c(2C_l \underline{Z}(0))} \end{aligned}$$

By Lemma 4, (2.5) and (2.6) we can construct a partition $(J_n)_{1 \leq n \leq N}$ of $[0, T']$ such that $\|u\|_{L_t^4 L_x^{12}(J_n \times \mathbb{R}^3)} = \frac{\epsilon}{g^{\frac{1}{4}}((2C_l)^n Z_0)}$, $1 \leq n < N$, $\|u\|_{L_t^4 L_x^{12}(J_N \times \mathbb{R}^3)} \leq \frac{\epsilon}{g^{\frac{1}{4}}((2C_l)^N Z_0)}$, $Z(t) \leq (2C_l)^n Z(0)$ for $t \in J_1 \cup \dots \cup J_n$ and

$$(2.7) \quad \frac{N^{\frac{1}{2}}}{\log^c(2C_l \underline{Z}(0))} \leq \max \left(C_{L,0}, (C_{L,1} E g(2M_0))^{C_{L,2} (E^{\frac{193}{4} + g} \frac{225}{8} + (2M_0))} \right)$$

Since $c < \frac{8}{225}$ we have by (1.24)

$$(2.8) \quad \begin{aligned} \log N &\lesssim \log^c(2C_l \underline{Z}(0)) + \log(C_{L,0}) + C_{L,2} E^{\frac{193}{4} + g} \log^{\frac{225c}{8}} + \log(10 + 4M_0^2) \log(C_{L,1} E \log^c \log(10 + 4M_0^2)) \\ &\leq \log \left(\frac{\log(\frac{M_0}{\underline{Z}(0)})}{\log(2C_l)} \right) \end{aligned}$$

if $M_0 = M_0 \left(\|u_0\|_{\tilde{H}^2(\mathbb{R}^3)}, \|u_1\|_{H^1(\mathbb{R})} \right)$ is large enough. To prove the last inequality in (2.8) it is enough, by using (1.24), to notice that $\lim_{M_0 \rightarrow \infty} f(M_0) = 0$ with

¹ such as the product rule $\log(a_1 a_2) = \log(a_1) + \log(a_2)$

$$(2.9) \quad f(M_0) := \frac{\log^c(2C_l \underline{Z}(0)) + \log(C_{L,0}) + C_{L,2} E^{\frac{193}{4}} + \log \frac{225c}{8} + \log(10+4M_0^2) \log(C_{L,1} E \log^c \log(10+4M_0^2))}{\log\left(\frac{\log\left(\frac{M_0}{\underline{Z}(0)}\right)}{\log(2C_l)}\right)}$$

Therefore we conclude that

$$(2.10) \quad \sup_{t \in [0, T']} \|(u(t), \partial_t u(t))\|_{H^2(\mathbb{R}^3) \times H^1(\mathbb{R}^3)} \leq (2C_l)^N Z(0) \leq M_0$$

2.1. Proof of Lemma 4. By the Strichartz estimates (1.16), the Sobolev embeddings (1.22) and (1.23) and the elementary estimate $|u^5 \nabla(g(u))| \lesssim |u^4 \nabla u g(u)|$

$$(2.11) \quad \begin{aligned} Z(t) &\lesssim Z(t_1) + \|u^5 g(u)\|_{L_t^1 L_x^2([t_1, t] \times \mathbb{R}^3)} + \|u^4 \nabla u g(u)\|_{L_t^1 L_x^2([t_1, t] \times \mathbb{R}^3)} + \|u^5 \nabla(g(u))\|_{L_t^1 L_x^2([t_1, t] \times \mathbb{R}^3)} \\ &\lesssim Z(t_1) + \|u^5 g(u)\|_{L_t^1 L_x^2([t_1, t] \times \mathbb{R}^3)} + \|u^4 \nabla u g(u)\|_{L_t^1 L_x^2([t_1, t] \times \mathbb{R}^3)} \\ &\lesssim Z(t_1) + \|u\|_{L_t^4 L_x^{12}([t_1, t] \times \mathbb{R}^3)}^4 \|u\|_{L_t^\infty L_x^6([t_1, t] \times \mathbb{R}^3)} \|g(u)\|_{L_t^\infty L_x^\infty([t_1, t] \times \mathbb{R}^3)} \\ &\quad + \|u\|_{L_t^4 L_x^{12}([t_1, t] \times \mathbb{R}^3)}^4 \|\nabla u\|_{L_t^\infty L_x^6([t_1, t] \times \mathbb{R}^3)} \|g(u)\|_{L_t^\infty L_x^\infty([t_1, t] \times \mathbb{R}^3)} \\ &\lesssim Z(t_1) + \|u\|_{L_t^4 L_x^{12}([t_1, t] \times \mathbb{R}^3)}^4 Z(t) g(Z(t)) \end{aligned}$$

Let C_l be the constant determined by the last inequality in (2.11). From (2.1), (2.11) and a continuity argument we have (2.2).

3. PROOF OF PROPOSITION 3

The proof relies upon five lemmas that are proved in the next sections.

Lemma 5. "Long time estimate if energy small" Let $J = [t_1, t_2]$ be a time interval. Let u be a classical solution of (1.1). Assume that (1.31) holds. If

$$(3.1) \quad E << \frac{1}{g^{\frac{1}{2}}(M)}$$

then

$$(3.2) \quad \|u\|_{L_t^4 L_x^{12}(J \times \mathbb{R}^3)} \lesssim 1$$

Lemma 6. "If $\|u\|_{L_t^4 L_x^{12}(J \times \mathbb{R}^3)}$ non negligible then existence of a mass concentration bubble and size of J bounded from below" Let u be a classical solution of (1.1). Let J be a time interval. Assume that (1.31) holds. Let η be a positive number such that

$$(3.3) \quad \eta \leq \frac{E^{\frac{1}{12}}}{g^{\frac{1}{24}}(M)}$$

If $\|u\|_{L_t^4 L_x^{12}(J \times \mathbb{R}^3)} \geq \eta$ then

$$(3.4) \quad \|u\|_{L_t^\infty L_x^6(J \times \mathbb{R}^3)} \gtrsim \eta^{2+} E^{-(\frac{1}{2}+)}$$

Moreover there exist a point $x_0 \in \mathbb{R}^3$, a time $t_0 \in J$ and a positive number r such that we have the mass concentration estimate in the ball $B(x_0, r)$

$$(3.5) \quad \int_{B(x_0, r)} |u(t_0, y)|^2 dy \gtrsim \eta^{6+} E^{-(2+)} r^2$$

and the following lower bound on the size of J

$$(3.6) \quad |J| \gtrsim \eta^4 E^{-\frac{2}{3}} r$$

Lemma 7. "Potential energy decay estimate" Let u be a classical solution of (1.1). Let $[a, b]$ be an interval. Then we have the potential energy decay estimate

$$(3.7) \quad \int_{|x| \leq b} F(u(b, x)) dx \lesssim \frac{a}{b} \left(e(a) + e^{\frac{1}{3}}(a) \right) + e(b) - e(a) + (e(b) - e(a))^{\frac{1}{3}}$$

Lemma 8. " $L_t^4 L_x^{12}$ norm of u is small on a large truncation of the forward light cone" Let $J = [t_1, t_2]$ be an interval. Let u be a classical solution of (1.1). Assume that (1.31) holds. Let η be a positive number such that

$$(3.8) \quad \eta \ll \min \left(E^{\frac{1}{4}}, E^{\frac{5}{18}}, \frac{E^{\frac{1}{12}}}{g^{\frac{1}{24}}(M)} \right)$$

Assume also that there exists $C_2 \gg 1$ such that

$$(3.9) \quad \left[t_1, (C_2 E^{10+} \eta^{-(36+)})^{4C_2 E^{10+} \eta^{-(36+)}} t_1 \right] \subset J$$

Then there exists a subinterval $J' = [t'_1, t'_2]$ such that $\left| \frac{t'_2}{t'_1} \right| \sim E^{10+} \eta^{-(36+)}$ and

$$(3.10) \quad \|u\|_{L_t^4 L_x^{12}(\Gamma_+(J'))} \leq \eta$$

Lemma 9. "Asymptotic stability" Let $J = [t_1, t_2]$ be a time interval. Let $J' = [t'_1, t'_2] \subset J$ and let $t \in J/J'$. Let u be a classical solution of (1.1). Assume that (1.31) holds. Then

$$(3.11) \quad \|u_{l, t'_2}(t) - u_{l, t'_1}(t)\|_{L_x^\infty(\mathbb{R}^3)} \lesssim \frac{E^{\frac{5}{6}} g^{\frac{1}{6}}(M)}{\text{dist}^{\frac{1}{2}}(t, J')}$$

We are ready to prove Proposition 3. We assume that we have an *a priori* bound M of the $L_t^\infty \dot{H}^2(J \times \mathbb{R}^3)$ norm of the solution u . There are two steps

- If $E \ll \frac{1}{g^{\frac{1}{2}}(M)}$ then we know from Lemma 5 that (1.32) holds.

- Therefore we assume that the energy is large, i.e

$$(3.12) \quad E \gtrsim \frac{1}{g^{\frac{1}{2}}(M)}$$

We can assume without loss of generality that

$$(3.13) \quad \|u\|_{L_t^4 L_x^{12}(J \times \mathbb{R}^3)} \geq \frac{E^{\frac{1}{12}}}{g^{\frac{1}{24}}(M)}$$

From (3.13) we can partition J into subintervals $J_1, \dots, J_l$ such that for $i = 1, \dots, l-1$

$$(3.14) \quad \|u\|_{L_t^4 L_x^{12}(J_i \times \mathbb{R}^3)} = \frac{E^{\frac{1}{12}}}{g^{\frac{1}{24}}(M)}$$

and

$$(3.15) \quad \|u\|_{L_t^4 L_x^{12}(J_l \times \mathbb{R}^3)} \leq \frac{E^{\frac{1}{12}}}{g^{\frac{1}{24}}(M)}$$

Before moving forward we say that an interval J_i is *exceptional* if ²

$$(3.16) \quad \|u_{l,t_1}\|_{L_t^4 L_x^{12}(J_i \times \mathbb{R}^3)} + \|u_{l,t_2}\|_{L_t^4 L_x^{12}(J_i \times \mathbb{R}^3)} \geq \frac{1}{(C_3 E g(M))^{C_4 \left(E^{\frac{193}{4} + g^{\frac{225}{8}} + (M) \right)}}$$

for some $C_3 \gg 1$, $C_4 \gg 1$ to be chosen later. Otherwise J_i is *unexceptional*. Let $\mathcal{E}$ denote the set of J_i 's that are exceptional and let $\overline{\mathcal{E}^c}$ denote the set of nonempty sequences of consecutive unexceptional intervals J_i . By (1.16), (3.12) and (3.16)

$$(3.17) \quad \begin{aligned} \text{card}(\mathcal{E}) &\lesssim E^2 [O(Eg(M))]^{O\left(E^{\frac{193}{4} + g^{\frac{225}{8}} + (M)}\right)} \\ &\lesssim [O(Eg(M))]^{O\left(E^{\frac{193}{4} + g^{\frac{225}{8}} + (M)}\right)} \end{aligned}$$

Since $\text{card}(\overline{\mathcal{E}^c}) \lesssim \text{card}(\mathcal{E})$ we have

$$(3.18) \quad \|u\|_{L_t^4 L_x^{12}(J \times \mathbb{R}^3)}^4 \lesssim [O(Eg(M))]^{O\left(E^{\frac{193}{4} + g^{\frac{225}{8}} + (M)}\right)} \left(\frac{E^{\frac{1}{3}}}{g^{\frac{1}{6}}(M)} + \sup_{K \in \overline{\mathcal{E}^c}} \|u\|_{L_t^4 L_x^{12}(K \times \mathbb{R}^3)}^4 \right)$$

Let $K = J_{i_0} \cup \dots \cup J_{i_1}$ is a sequence of consecutive unexceptional intervals. If $N(K)$ is the number of J_i 's making K then by (3.12), (3.14), (3.15) and (3.18) we have

$$(3.19) \quad \|u\|_{L_t^4 L_x^{12}(J \times \mathbb{R}^3)} \lesssim (\sup_{K \in \overline{\mathcal{E}^c}} N(K)) [O(Eg(M))]^{O\left(E^{\frac{193}{4} + g^{\frac{225}{8}} + (M)}\right)}$$

Therefore it suffices to estimate $N(K)$ for every $K = J_{i_0} \cup \dots \cup J_{i_1}$. We will do that by first determining a lower bound for the size of the elements J_i 's and then by estimating the size of K . By (3.12), (3.14), (3.15) and Lemma 6 there exists for $i \in [i_0, \dots, i_1]$ a $(t_i, r_i, x_i) \in (J_i \times (0, \infty) \times \mathbb{R}^3)$ such that

²The numbers $\frac{193}{4}$ and $\frac{225}{8}$ will play an important role in (3.45)

$$(3.20) \quad \frac{1}{r_i^2} \int_{B(x_i, r_i)} |u(t_i, y)|^2 dy \gtrsim \frac{E^{-(\frac{3}{2}+)}}{g^{\frac{5}{4}+}(M)}$$

and

$$(3.21) \quad |J_i| \gtrsim \frac{E^{-\frac{1}{3}} r_i}{g^{\frac{5}{6}}(M)}$$

Let $k \in [i_0, \dots, i_1]$ such that $r_k = \min_{i \in [i_0, i_1]} r_i$, let $f(t, r, x) := \frac{1}{r^2} \int_{B(x, r)} |u(t, y)|^2 dy$, let C_5 be the constant determined by (3.20). Let $r_0 = r_0(M)$ such that $r_0 M^2 = \frac{C_5 E^{-(\frac{3}{2}+)}}{4g^{\frac{5}{4}+}(M)}$. Since $f(t, r, x) \leq r M^2$ we have $f(t, r_0, x) \leq \frac{C_5 E^{-(\frac{3}{2}+)}}{4g^{\frac{5}{4}+}(M)}$. The set $A := \{(t, r, x) : t \in K, r_0 \leq r \leq r_k, x \in \mathbb{R}^3\}$ is connected. Therefore its image is connected by f and there exists $(\tilde{t}, \tilde{r}, \tilde{x}) \in K \times [r_0, r_k] \times \mathbb{R}^3$ such that $f(\tilde{t}, \tilde{r}, \tilde{x}) = \frac{C_5 E^{-(\frac{3}{2}+)}}{2g^{\frac{5}{4}+}(M)}$. In other words we have the following mass concentration

$$(3.22) \quad \frac{1}{\tilde{r}^2} \int_{B(\tilde{x}, \tilde{r})} u^2(\tilde{t}, x) dx = \frac{C_5 E^{-(\frac{3}{2}+)}}{2g^{\frac{5}{4}+}(M)}$$

Moreover we have the useful lower bound for the size of J_i , $i_0 \leq i \leq i_1$ ³

$$(3.23) \quad |J_i| \gtrsim \tilde{r} \frac{E^{-\frac{1}{3}}}{g^{\frac{5}{6}}(M)}$$

At this point we need to use the following lemma that we will prove in the next subsection

Lemma 10. *Let K be a sequence of unexceptional intervals. Assume that there exists $(\bar{t}, \bar{x}, \bar{r}) \in K \times \mathbb{R}^3 \times (0, \infty)$ such that*

$$(3.24) \quad \frac{1}{\bar{r}^2} \int_{B(\bar{x}, \bar{r})} u^2(\bar{t}, y) dy \gtrsim E^{-(\frac{3}{2}+)} g^{\frac{5}{4}+}(M)$$

Then there exist two constants $C_6 \gg 1$, $C_7 \gg 1$ such that

$$(3.25) \quad |K| \leq (C_6 E g(M))^{C_7 E^{\frac{193}{4}+} g^{\frac{225}{8}+}(M)} \bar{r}$$

Lemma 10 gives information about the size of K . We postpone the proof of this lemma to the next subsection. If we combine it to (3.23) we can estimate $N(K)$. More precisely by Lemma 10, (3.23) and (3.12) we have

³ Notice that we have the lower bound $\tilde{r} \geq \frac{C_5 E^{-(\frac{3}{2}+)}}{4M^2 g^{\frac{5}{4}+}(M)}$. One might think that the presence of $\tilde{r}$ in (3.23) is annoying since this lower bound is crude. However we will see that $\tilde{r}$ disappears at the end of the process: see (3.26). Therefore a sharp lower bound is not required.

$$\begin{aligned}
(3.26) \quad N(K) &\lesssim \frac{(C_6 E g(M))^{C_7 E^{\frac{193}{4} + g^{\frac{225}{8} + (M)}} \bar{r}}}{\bar{r}^{\frac{E^{-\frac{1}{3}}}{g^{\frac{5}{8} + (M)}}}} \\
&\lesssim (O(E g(M)))^{O(E^{\frac{193}{4} + g^{\frac{225}{8} + (M)}})}
\end{aligned}$$

Plugging this upper bound of $N(K)$ into (3.19) we get (1.33).

3.1. Proof of Lemma 10. By using the space translation invariance of (1.1) we can reduce to the case $\bar{x} = 0$ ⁴. By using the time reversal invariance⁵ it suffices to estimate $|K \cap [\bar{t}, \infty)|$. By using the time translation invariance again⁶ we can assume that $\bar{t} = \bar{r}$ and therefore $\bar{r} \in K$. Let $K_+ := K \cap [\bar{r}, \infty)$. We are interested in estimating $|K_+|$. We would like to use Lemma 8. Therefore, we consider the set $\Gamma_+(K_+)$. We have

$$(3.27) \quad \frac{1}{\bar{r}^2} \int_{B(0, \bar{r})} |u(\bar{r}, y)|^2 dy \gtrsim \frac{E^{-(\frac{3}{2}+)}}{g^{\frac{5}{4} + (M)}}$$

Therefore by Proposition 2 and (3.27) we have

$$(3.28) \quad \int_{B(0, \bar{r})} |u(t, y)|^2 dy \gtrsim \frac{E^{-(\frac{3}{2}+) \bar{r}^2}}{g^{\frac{5}{4} + (M)}}$$

if $(t - \bar{r})E^{\frac{1}{2}} \leq \frac{c_0 E^{-(\frac{3}{4}+) \bar{r}}}{g^{\frac{5}{8} + (M)}}$ for some $c_0 \ll 1$. Therefore by Hölder there exists $0 < c_1 \ll 1$ small enough such that

$$\begin{aligned}
(3.29) \quad &\|u\|_{L_t^4 L_x^{12} \left(\Gamma_+ \left(\left[\bar{r}, \bar{r} + \frac{c_0 E^{-(\frac{5}{4}+) \bar{r}}}{g^{\frac{5}{8} + (M)}} \right] \right) \right)} \geq c_1 \frac{E^{-\frac{17}{16}}}{g^{\frac{25}{32} + (M)}} \\
&- \|u\|_{L_t^4 L_x^{12}(\Gamma_+(K_+))} \leq c_1 \frac{E^{-\frac{17}{16}}}{g^{\frac{25}{32} + (M)}}. \text{ In this case we get from (3.29)}
\end{aligned}$$

$$(3.30) \quad K_+ \subset \left[\bar{r}, \bar{r} + \frac{c_0 E^{-(\frac{5}{4}+) \bar{r}}}{g^{\frac{5}{8} + (M)}} \right]$$

and, using also (3.12), we get (3.25).

$$\begin{aligned}
&- \|u\|_{L_t^4 L_x^{12}(\Gamma_+(K_+))} \geq c_1 \frac{E^{-(\frac{17}{16}+)}}{g^{\frac{25}{32} + (M)}}. \text{ In the sequel we denote by } \tilde{\eta} \text{ the} \\
&\text{following number}
\end{aligned}$$

$$(3.31) \quad \tilde{\eta} := \frac{c_1}{4} \frac{E^{-(\frac{17}{16}+)}}{g^{\frac{25}{32} + (M)}}$$

⁴we consider the function $u_1(t, x) = u(t, x - \bar{x})$ and we abuse notation in the sequel by writing u_1 for u

⁵we consider the function $u_2(t, x) := u(2\bar{t} - t, x)$ and we abuse notation in the sequel by writing u_2 for u

⁶we consider the function $u_3(t, x) := u(t + (\bar{t} - \bar{r}), x)$ and we abuse notation in the sequel by writing u_3 for u

and we divide $\Gamma_+(K_+)$ into consecutive cone truncations $\Gamma_+(\tilde{J}_1), \dots, \Gamma_+(\tilde{J}_k)$ such that for $j = 1, \dots, k-1$

$$(3.32) \quad \|u\|_{L_t^4 L_x^{12}(\Gamma_+(\tilde{J}_j))} = \tilde{\eta}$$

and

$$(3.33) \quad \|u\|_{L_t^4 L_x^{12}(\Gamma_+(\tilde{J}_k))} \leq \tilde{\eta}$$

We get from (3.29)

$$(3.34) \quad \tilde{J}_1 \subset \left[\bar{r}, \bar{r} + \frac{c_0 E^-(\frac{5}{4}) \bar{r}}{g^{\frac{5}{8}+(M)}} \right]$$

Now we prove the following result before moving forward

Result 1. *If $j \in [1, \dots, k-1]$ we either have*

$$(3.35) \quad |\tilde{J}_{j+1}| \lesssim |\tilde{J}_j| \tilde{\eta}^{-4} E^{\frac{8}{3}} g^{\frac{1}{3}}(M)$$

or

$$(3.36) \quad |\tilde{J}_j| \geq (C_6 E g(M))^{C_7 E^{\frac{193}{4} + g^{\frac{225}{8} + (M)}} \bar{r}}$$

for some constants $C_6 \gg 1, C_7 \gg 1$.

Proof. We get from (1.21), (3.12) and (3.31)

$$(3.37) \quad \begin{aligned} \|u - u_{l,t_{j+1}}\|_{L_t^4 L_x^{12}(\Gamma_+(\tilde{J}_j))} &\lesssim \|u^5 g(u)\|_{L_t^1 L_x^2(\Gamma_+(\tilde{J}_j \cup J_{j+1}))} \\ &\lesssim \|u^4\|_{L_t^1 L_x^3(\Gamma_+(\tilde{J}_j \cup J_{j+1}))} \|u g^{\frac{1}{6}}(u)\|_{L_t^\infty L_x^6(\Gamma_+(\tilde{J}_j \cup J_{j+1}))} g^{\frac{5}{6}}(M) \\ &\lesssim \tilde{\eta}^4 E^{\frac{1}{6}} g^{\frac{5}{6}}(M) \\ &<< \tilde{\eta} \end{aligned}$$

with $J_j = [t_{j-1}, t_j]$. Therefore by (3.32) we have $\|u_{l,t_{j+1}}\|_{L_t^4 L_x^{12}(\Gamma_+(\tilde{J}_j))} \sim \tilde{\eta}$. This implies that

$$(3.38) \quad \|u_{l,t_{j+1}} - u_{l,t_2}\|_{L_t^4 L_x^{12}(\Gamma_+(\tilde{J}_j))} \gtrsim \tilde{\eta}.$$

or

$$(3.39) \quad \|u_{l,t_2}\|_{L_t^4 L_x^{12}(\Gamma_+(\tilde{J}_j))} \gtrsim \tilde{\eta}$$

– *Case 1:* $\|u_{l,t_{j+1}} - u_{l,t_2}\|_{L_t^4 L_x^{12}(\Gamma_+(\tilde{J}_i))} \gtrsim \tilde{\eta}$. By Lemma 9 and Hölder we have

$$(3.40) \quad \begin{aligned} \|u_{l,t_{j+1}} - u_{l,t_2}\|_{L_t^4 L_x^{12}(\Gamma_+(\tilde{J}_j))} &\lesssim |\tilde{J}_j|^{\frac{1}{4}} \|u_{l,t_{j+1}} - u_{l,t_2}\|_{L_t^\infty L_x^{12}(\Gamma_+(\tilde{J}_j))} \\ &\lesssim |\tilde{J}_j|^{\frac{1}{4}} \|u_{l,t_{j+1}} - u_{l,t_2}\|_{L_t^\infty L_x^\infty(\Gamma_+(\tilde{J}_j))}^{\frac{1}{2}} \|u_{l,t_{j+1}} - u_{l,t_2}\|_{L_t^\infty L_x^6(\Gamma_+(\tilde{J}_j))}^{\frac{1}{2}} \\ &\lesssim \frac{|\tilde{J}_j|^{\frac{1}{4}} E^{\frac{2}{3}} g^{\frac{1}{12}}(M)}{|\tilde{J}_{j+1}|^{\frac{1}{4}}} \end{aligned}$$

We get (3.35) from (3.38) and (3.40).

– *Case 2:* $\|u_{l,t_2}\|_{L_t^4 L_x^{12}(\Gamma_+(\tilde{J}_j))} \gtrsim \tilde{\eta}$. In this case $\|u_{l,t_2}\|_{L_t^4 L_x^{12}(\tilde{J}_j)} \gtrsim \tilde{\eta}$. Recall that K_+ is a subinterval of $K = J_{i_0} \cup \dots \cup J_{i_1}$, sequence of unexceptional intervals J_i , $i_0 \leq i \leq i_1$. Consequently there are at least $\sim \tilde{\eta} (C_3 E g(M))^{C_4 E^{\frac{193}{4}+g^{\frac{225}{8}+}(M)}}$ intervals J_j that cover $\tilde{J}_i$. Therefore we get (3.36) from (3.23) and (3.12). $\square$

Using Result 1 and Lemma 8 we can get an upper bound on the size $|K_+|$

Result 2. *We have*

$$(3.41) \quad |K_+| \leq (C_6 E g(M))^{C_7 (E^{\frac{193}{4}+g^{\frac{225}{8}+}(M)})} \bar{r}$$

Proof. Let $B := (C_6 E g(M))^{C_7 (E^{\frac{193}{4}+g^{\frac{225}{8}+}(M)})}$. Assume that (3.41) fails. Let $\tilde{J}_{j_1}$ be the first interval for which $|\tilde{J}_1 \cup \dots \cup \tilde{J}_{j_1}|$ exceeds $B\bar{r}$. Then $j_1 \neq 1$, $|\tilde{J}_{j_1}| \lesssim |\tilde{J}_{j_1-1}| \tilde{\eta}^{-4} E^{\frac{8}{3}} g^{\frac{1}{3}}(M)$ and we have

$$(3.42) \quad \frac{c_1 E^{-\frac{5}{4}} \bar{r}}{g^{\frac{8}{3}}(M)} + T_2 - T_1 + (T_2 - T_1) \tilde{\eta}^{-4} E^{\frac{8}{3}} g^{\frac{1}{3}}(M) \gtrsim |\tilde{J}_1| + \dots + |\tilde{J}_{j_1}| \geq B\bar{r}$$

if $[T_1, T_2] := \tilde{J}_2 \cup \dots \cup \tilde{J}_{j_1-1}$. Therefore by (3.12) and (3.42) we have

$$(3.43) \quad T_2 - T_1 \gtrsim \frac{\tilde{\eta}^4 E^{-\frac{8}{3}} B\bar{r}}{g^{\frac{1}{3}}(M)}$$

Moreover $T_1 \leq \bar{r} + \frac{c_1 E^{-\frac{5}{4}} \bar{r}}{g^{\frac{8}{3}}(M)}$. Therefore by (3.12) we have

$$(3.44) \quad T_1 = O(\bar{r})$$

By (3.43) and (3.44) we have

$$(3.45) \quad \frac{T_2}{T_1} \geq \left(C_2 E^{10+} \left(\frac{\tilde{\eta}}{4} \right)^{-(36+)} \right)^{4C_2 E^{10+} \left(\frac{\tilde{\eta}}{4} \right)^{-(36+)}}$$

with C_2 defined in Lemma 8, provided that $C_6, C_7 \gg \max(c_1, C_2)$. Therefore we can apply Lemma 8 and find a subinterval $[t'_1, t'_2] \subset \tilde{J}_2 \cup \dots \cup \tilde{J}_{j_1-1}$ with $\left| \frac{t'_2}{t'_1} \right| \sim E^{10+} \tilde{\eta}^{-(36+)}$ and $\|u\|_{L_t^4 L_x^{12}([t'_1, t'_2])} \leq \frac{\tilde{\eta}}{4}$. This means that $[t'_1, t'_2] \subset [T_1, T_2]$ is covered by at most two consecutive intervals. It is convenient to introduce $[t'_1, t'_2]_g$, the geometric mean of t'_1 and t'_2 . We have $[t'_1, t'_2]_g \sim \tilde{\eta}^{-18} E^5 t'_1$. There are two cases

– *Case 1:* $[t'_1, t'_2]$ is covered by one interval $\tilde{J}_{\bar{j}} = [a_{\bar{j}}, b_{\bar{j}}]$, $2 \leq \bar{j} \leq j_1 - 1$. Then $|\tilde{J}_{\bar{j}}| \gtrsim \tilde{\eta}^{-(36+)} E^{10+} t'_1$ and $|\tilde{J}_{\bar{j}-1}| \leq t'_1$. Therefore $|\tilde{J}_{\bar{j}}| \gtrsim \tilde{\eta}^{-(36+)} E^{10+} |\tilde{J}_{\bar{j}-1}|$. Contradiction with (3.12) and (3.35).

- *Case 2:* $[t'_1, t'_2]$ is covered by two intervals $\tilde{J}_{\bar{j}} = [a_{\bar{j}}, b_{\bar{j}}]$ and $\tilde{J}_{\bar{j}+1} = [a_{\bar{j}+1}, b_{\bar{j}+1}]$ for some $2 \leq \bar{j} \leq j_1 - 2$. Then there are two subcases
 - * *Case 2.a:* $b_{\bar{j}} \leq [t'_1, t'_2]_g$. In this case $|\tilde{J}_{\bar{j}+1}| \gtrsim \tilde{\eta}^{-(36+)} E^{10+t'_1}$ and $|\tilde{J}_{\bar{j}}| \leq \tilde{\eta}^{-(18+)} E^{5+t'_1}$. Therefore by (3.12) we have $|\tilde{J}_{\bar{j}+1}| \gtrsim \tilde{\eta}^{-(18+)} E^{5+|\tilde{J}_{\bar{j}}|}$. Contradiction with (3.12) and (3.35).
 - * *Case 2.b:* $b_{\bar{j}} \geq [t'_1, t'_2]_g$. In this case by (3.12) $|\tilde{J}_{\bar{j}}| \gtrsim \tilde{\eta}^{-(18+)} E^{5+t'_1}$ and $|\tilde{J}_{\bar{j}-1}| \leq t'_1$. Therefore $|\tilde{J}_{\bar{j}}| \gtrsim \tilde{\eta}^{-(18+)} E^{5+|\tilde{J}_{\bar{j}-1}|}$. Contradiction with (3.12) and (3.35).

□

Remark 3.1. *It seems likely that we can find a better upper bound for $|K_+|$ than (3.41) by exploiting Lemma 9 in a better way. For instance we can consider k successive time intervals $\tilde{J}_{j+1}, \dots, \tilde{J}_{j+k}$, $k > 1$ and prove an estimate like*

$$(3.46) \quad |\tilde{J}_{j+1}| + \dots |\tilde{J}_{j+k}| \lesssim |\tilde{J}_{\bar{j}}| \tilde{\eta}^{-4} E^{\frac{8}{5}} g^{\frac{1}{3}}(M)$$

This estimate is stronger than (3.35). We can probably find a smaller B such that (3.45) holds with $\tilde{\eta}$ substituted for something like $k\tilde{\eta}$ and, by modifying the argument above, find a contradiction with (3.46). At the end of the process we can probably prove global existence of smooth solutions to (1.1) for $0 < c < c_0$, with $c_0 > \frac{8}{225}$ to be determined. We will not pursue these matters.

4. PROOF OF LEMMA 5

Applying the Strichartz estimates and Hölder inequality

$$(4.1) \quad \begin{aligned} \|u\|_{L_t^4 L_x^{12}(J \times \mathbb{R}^3)} &\lesssim E^{\frac{1}{2}} + \|u^4\|_{L_t^1 L_x^2(J \times \mathbb{R}^3)} \|ug^{\frac{1}{6}}(u)\|_{L_t^\infty L_x^6(J \times \mathbb{R}^3)} \|g^{\frac{5}{6}}(u)\|_{L_t^\infty L_x^\infty(J \times \mathbb{R}^3)} \\ &\lesssim E^{\frac{1}{2}} + E^{\frac{1}{6}} g^{\frac{5}{6}}(M) \|u\|_{L_t^4 L_x^{12}(J \times \mathbb{R}^3)}^4 \end{aligned}$$

Hence (3.2) by (3.1) and a continuity argument.

5. PROOF OF LEMMA 6

Let $J' = [t'_1, t'_2] \subset J$ be such that $\|u\|_{L_t^4 L_x^{12}(J' \times \mathbb{R}^3)} = \eta$. Then by (1.22) and (3.3)

$$(5.1) \quad \begin{aligned} \|f(u)\|_{L_t^1 L_x^2(J' \times \mathbb{R}^3)} &\lesssim \|ug^{\frac{1}{6}}(u)\|_{L_t^\infty L_x^6(J' \times \mathbb{R}^3)} \|u\|_{L_t^4 L_x^{12}(J' \times \mathbb{R}^3)}^4 \|g^{\frac{5}{6}}(u)\|_{L_t^\infty L_x^\infty(J' \times \mathbb{R}^3)} \\ &\lesssim E^{\frac{1}{6}} \eta^4 g^{\frac{5}{6}}(M) \\ &\lesssim E^{\frac{1}{2}} \end{aligned}$$

It is slightly unfortunate that $(2, \infty)$ is not wave admissible. Therefore we consider the admissible pair $\left(2 + \epsilon, \frac{6(2+\epsilon)}{\epsilon}\right)$ with $\epsilon \ll 1$. By the Strichartz estimates and (5.1)

$$\begin{aligned}
(5.2) \quad \|u\|_{L_t^{2+\epsilon} L_x^{\frac{6(2+\epsilon)}{\epsilon}}(J' \times \mathbb{R}^3)} &\lesssim \|\nabla u(t'_1)\|_{L^2(\mathbb{R}^3)} + \|u(t'_1)\|_{L^2(\mathbb{R}^3)} + \|f(u)\|_{L_t^1 L_x^2(J' \times \mathbb{R}^3)} \\
&\lesssim E^{\frac{1}{2}}
\end{aligned}$$

Moreover let N be a frequency to be chosen later. By Bernstein inequality and (1.7) we have

$$\begin{aligned}
(5.3) \quad \|P_{<N} u\|_{L_t^4 L_x^{12}(J' \times \mathbb{R}^3)} &\lesssim N^{\frac{1}{4}} |J'|^{\frac{1}{4}} \|u\|_{L_t^\infty L_x^6(J' \times \mathbb{R}^3)} \\
&\lesssim N^{\frac{1}{4}} |J'|^{\frac{1}{4}} E^{\frac{1}{6}}
\end{aligned}$$

Therefore

$$(5.4) \quad \|P_{<N} u\|_{L_t^4 L_x^{12}(J' \times \mathbb{R}^3)} \lesssim |J'|^{\frac{1}{4}} N^{\frac{1}{4}} E^{\frac{1}{6}}$$

Let $c_2 < 1$. Then if $N = c_2^4 \frac{\eta^4}{|J'| E^{\frac{2}{3}}}$ we have

$$(5.5) \quad \|P_{\geq N} u\|_{L_t^4 L_x^{12}(J' \times \mathbb{R}^3)} \gtrsim \eta$$

and

$$(5.6) \quad \|u\|_{L_t^4 L_x^{12}(J' \times \mathbb{R}^3)} \sim \|P_{\geq N} u\|_{L_t^4 L_x^{12}(J' \times \mathbb{R}^3)}$$

By (5.2), (5.5) and (5.6) we have

$$\begin{aligned}
(5.7) \quad \eta &\sim \|P_{\geq N} u\|_{L_t^4 L_x^{12}(J' \times \mathbb{R}^3)} \\
&\lesssim \|P_{\geq N} u\|_{L_t^{2+\epsilon} L_x^{\frac{6(2+\epsilon)}{\epsilon}}(J' \times \mathbb{R}^3)}^{\frac{2+\epsilon}{4}} \|P_{\geq N} u\|_{L_t^\infty L_x^6(J' \times \mathbb{R}^3)}^{1-\frac{2+\epsilon}{4}} \\
&\lesssim E^{\frac{2+\epsilon}{8}} \|P_{\geq N} u\|_{L_t^\infty L_x^6(J' \times \mathbb{R}^3)}^{1-\frac{2+\epsilon}{4}}
\end{aligned}$$

Therefore we conclude that $\|P_{\geq N} u\|_{L_t^\infty L_x^6(J' \times \mathbb{R}^3)} \gtrsim \eta^{2+} E^{-(\frac{1}{2}+)}$. Applying Proposition 1 we get (3.5).

6. PROOF OF LEMMA 7

Bahouri and Gerard (see [1], p171) used arguments from Grillakis [4, 5] and Shatah-Struwe [12] to derive an a priori estimate of the solution u to the 3D quintic defocusing wave equation, i.e $\partial_{tt} u - \Delta u + u^5 = 0$. More precisely they were able to prove

$$(6.1) \quad \int_{|x| \leq b} |u(b, x)|^6 dx \lesssim \frac{a}{b} \left(\tilde{e}(a) + \tilde{e}^{\frac{1}{3}}(a) \right) + \tilde{e}(b) - \tilde{e}(a) + (\tilde{e}(b) - \tilde{e}(a))^{\frac{1}{3}}$$

with

$$(6.2) \quad \tilde{e}(t) := \frac{1}{2} \int_{|x| \leq t} (\partial_t u)^2 dx + \frac{1}{2} \int_{|x| \leq t} |\nabla u|^2 dx + \frac{1}{6} \int_{|x| \leq t} u^6 dx$$

Since we apply their ideas to the potential f we just sketch the proof. Given the cone $\Gamma_+([a, b])$ we denote by $\partial\Gamma_+([a, b])$ the mantle of the cone $\Gamma_+([a, b])$ i.e

$$(6.3) \quad \partial\Gamma_+([a, b]) := \left\{ (t', x) \in [a, b] \times \mathbb{R}^3, t = |x| \right\}$$

The local energy identity

$$(6.4) \quad e(b) - e(a) = \frac{1}{2\sqrt{2}} \int_{\partial\Gamma_+([a, b])} \left| \frac{x\partial_t u}{t} + \nabla u \right|^2 + \frac{1}{\sqrt{2}} \int_{\partial\Gamma_+([a, b])} F(u)$$

results from the integration of the identity $\partial_t u (\partial_{tt} u - \Delta u + f(u)) = 0$ on the cone $\Gamma_+([a, b])$. We have [11]

$$(6.5) \quad \begin{aligned} & \partial_t \left(\frac{t}{2} (\partial_t u)^2 + \frac{t}{2} |\nabla u|^2 + (x \cdot \nabla u) \partial_t u + tF(u) + u \partial_t u \right) \\ & - \operatorname{div} \left(t \nabla u \partial_t u + (x \cdot \nabla u) \nabla u - \frac{|\nabla u|^2 x}{2} + \frac{(\partial_t u)^2 x}{2} - xF(u) + u \nabla u \right) + u f(u) - 4F(u) \\ & = 0 \end{aligned}$$

Integrating this identity on $\Gamma_+([a, b])$ we have

$$(6.6) \quad X(b) - X(a) + Y(a, b) = \int_{\Gamma_+([a, b])} 4F(u) - u f(u)$$

with

$$(6.7) \quad X(t) := \int_{|x| \leq t} \frac{t}{2} (\partial_t u)^2 + \frac{t}{2} |\nabla u|^2 + (x \cdot \nabla u) \partial_t u + tF(u) + u \partial_t u$$

and

$$(6.8) \quad Y(a, b) := -\frac{1}{\sqrt{2}} \int_{\partial\Gamma_+([a, b])} \left[\frac{t}{2} (\partial_t u)^2 + \frac{t}{2} |\nabla u|^2 + (x \cdot \nabla u) \partial_t u + tF(u) + u \partial_t u + t \frac{\nabla u \cdot x}{|x|} \partial_t u + \frac{|x \cdot \nabla u|^2}{|x|} \right. \\ \left. - \frac{|\nabla u|^2 |x|}{2} + \frac{(\partial_t u)^2 |x|}{2} - |x| F(u) + u \frac{\nabla u \cdot x}{|x|} \right]$$

In fact (see [12]) we have

$$(6.9) \quad X(t) = \int_{|x| \leq t} t \left[\frac{1}{2} (\partial_t u)^2 + \frac{1}{2} \left| \nabla u + \frac{ux}{|x|^2} \right|^2 \right] + \partial_t u (x \cdot \nabla u + u) + tF(u) - \int_{|x|=t} \frac{u^2}{2}$$

Since $t = |x|$ on $\partial\Gamma_+([a, b])$ we have

$$(6.10) \quad Y(a, b) = -\frac{1}{\sqrt{2}} \int_{\partial\Gamma_+([a, b])} |x| (\partial_t u)^2 + 2(x \cdot \nabla u) \partial_t u + u \partial_t u + \frac{(x \cdot \nabla u)^2}{|x|} + u \frac{\nabla u \cdot x}{|x|}$$

and after some computations (see [12])

$$(6.11) \quad Y(a, b) = -\frac{1}{\sqrt{2}} \int_{\partial\Gamma_+([a, b])} \frac{1}{t} (t \partial_t u + (\nabla u \cdot x) + u)^2 + \int_{|x|=b} \frac{u^2}{2} - \int_{|x|=a} \frac{u^2}{2}$$

Therefore if

$$(6.12) \quad H(t) := \int_{|x| \leq t} t \left[\frac{1}{2} (\partial_t u)^2 + \frac{1}{2} \left| \nabla u + \frac{ux}{|x|^2} \right|^2 \right] + \partial_t u (x \cdot \nabla u + u) + tF(u)$$

then

$$(6.13) \quad H(b) - H(a) = \frac{1}{\sqrt{2}} \int_{\partial\Gamma_+([a,b])} \frac{1}{t} (t\partial_t u + \nabla u \cdot x + u)^2 + \int_{\Gamma_+([a,b])} 4F(u) - uf(u)$$

We estimate $H(t)$, following [1]. We have

$$(6.14) \quad \begin{aligned} |\partial_t u(x \cdot \nabla u + u)| &\leq \frac{t}{2} \left((\partial_t u)^2 + \left| \nabla u + \frac{ux}{|x|^2} \right|^2 \right) \\ &\lesssim t \left((\partial_t u)^2 + |\nabla u|^2 + \frac{u^2}{|x|^2} \right) \end{aligned}$$

Therefore by (6.14), Hölder inequality and (1.7)

$$(6.15) \quad \begin{aligned} H(t) &\lesssim t \left(e(t) + \int_{|x| \leq t} \frac{u^2}{|x|^2} \right) \\ &\lesssim t \left(e(t) + \left(\int_{|x| \leq t} u^6 \right)^{\frac{1}{3}} \right) \\ &\lesssim t \left(e(t) + e^{\frac{1}{3}}(t) \right) \end{aligned}$$

Moreover by (6.4), Hölder inequality and (1.7)

$$(6.16) \quad \begin{aligned} \frac{1}{\sqrt{2}} \int_{\partial\Gamma_+([a,b])} \frac{1}{t} (t\partial_t u + \nabla u \cdot x + u)^2 &\lesssim \frac{b}{2\sqrt{2}} \int_{\partial\Gamma_+([a,b])} \left(\frac{\nabla u \cdot x}{t} + \partial_t u \right)^2 + \frac{1}{2\sqrt{2}} \int_{\partial\Gamma_+([a,b])} \frac{u^2}{t^2} \\ &\lesssim b \int_{\partial\Gamma_+([a,b])} \left| \frac{x}{t} \partial_t u + \nabla u \right|^2 + \frac{1}{2\sqrt{2}} \left(\int_{\partial\Gamma_+([a,b])} u^6 \right)^{\frac{1}{3}} \\ &\lesssim b \left((e(b) - e(a)) + (e(b) - e(a))^{\frac{1}{3}} \right) \end{aligned}$$

We get from (1.7)

$$(6.17) \quad 4F(u) - uf(u) \leq 0$$

By (6.13), (6.15), (6.16) and (6.17) we have

$$(6.18) \quad \begin{aligned} \int_{|x| \leq b} F(u) &\lesssim \frac{H(b)}{b} \\ &\lesssim \frac{H(a) + \frac{1}{\sqrt{2}} \int_{\partial\Gamma_+([a,b])} \frac{1}{t} (t\partial_t u + \nabla u \cdot x + u)^2}{b} \\ &\lesssim \frac{a}{b} \left(e(a) + e^{\frac{1}{3}}(a) \right) + e(b) - e(a) + (e(b) - e(a))^{\frac{1}{3}} \end{aligned}$$

7. PROOF OF LEMMA 8

The proof relies upon two results that we prove in the subsections.

Result 3. *Let u be a classical solution of (1.1). Assume that (1.31) holds. Let η be a positive number such that (3.3) holds. If $\|u\|_{L_t^4 L_x^{12}(\Gamma_+(J))} \geq \eta$ then*

$$(7.1) \quad \|u\|_{L_t^\infty L_x^6(\Gamma_+(J))} \gtrsim \eta^{2+} E^{-(\frac{1}{2}+)}$$

Result 4. *Let u be a smooth solution to (1.1). Assume that (1.31) holds. Let η be a positive number such that*

$$(7.2) \quad \eta \leq \min \left(1, E^{\frac{1}{18}} \right)$$

Let $J = [t_1, t_2]$ be an interval such that $\left[t_1, t_1 (E\eta^{-18})^{4E\eta^{-18}} \right] \subset J$. Then there exists a subinterval $J' = [t'_1, t'_2]$ such that $\left| \frac{t'_2}{t'_1} \right| = E\eta^{-18}$ and

$$(7.3) \quad \|u\|_{L_t^\infty L_x^6(\Gamma_+(J'))} \lesssim \eta$$

Let C_9 be the constant determined by $\gtrsim$ in (7.1). Let C_{10} be the constant determined by $\lesssim$ in (7.3). We get from (3.9)

$$(7.4) \quad \left[t_1, t_1 \left(E \left(\frac{C_9 \eta^{2+E^-(\frac{1}{2})+}}{2C_{10}} \right)^{-18} \right)^{4E \left(\frac{C_9 \eta^{2+E^-(\frac{1}{2})+}}{2C_{10}} \right)^{-18}} \right] \subset \left[t_1, C_2 (E^{10+} \eta^{-(36+)})^{4C_2 E^{10+} \eta^{-(36+)}} t_1 \right] \subset J$$

if $C_2 \gg \max(C_9, C_{10})$. Therefore, since $\frac{C_9 \eta^{2+E^-(\frac{1}{2})+}}{2C_{10}}$ satisfies (7.2) by (3.8), we can use Result 4 and show that there exists a subinterval $J' = [t'_1, t'_2]$ such that $\left| \frac{t'_2}{t'_1} \right| \sim E^{10+} \eta^{-(36+)}$ and

$$(7.5) \quad \begin{aligned} \|u\|_{L_t^\infty L_x^6(\Gamma_+(J'))} &\leq \frac{C_9 \eta^{2+E^-(\frac{1}{2})+} C_{10}}{2C_{10}} \\ &\leq C_9 \frac{\eta^{2+E^-(\frac{1}{2})+}}{2} \end{aligned}$$

Now we claim that $\|u\|_{L_t^4 L_x^{12}(\Gamma_+(J'))} \leq \eta$. If not by (3.8) and Result 3 we have

$$(7.6) \quad \|u\|_{L_t^\infty L_x^6(\Gamma_+(J'))} \geq C_9 \eta^{2+E^-(\frac{1}{2})+}$$

Contradiction with (7.5).

7.1. Proof of Result 3. We substitute J' for $\Gamma_+(J')$ in (5.1) to get

$$(7.7) \quad \|f(u)\|_{L_t^1 L_x^2(\Gamma_+(J'))} \lesssim E^{\frac{1}{2}}$$

By the Strichartz estimates (1.20) on the truncated cone $\Gamma_+(J')$ we have

$$(7.8) \quad \|u\|_{L_t^{2+\epsilon} L_x^{\frac{6(2+\epsilon)}{\epsilon}}(\Gamma_+(J'))} \lesssim E^{\frac{1}{2}}$$

after following similar steps to prove (5.2). Therefore

$$\begin{aligned}
(7.9) \quad \eta &= \|u\|_{L_t^4 L_x^{12}(\Gamma_+(J))} \\
&\lesssim \|u\|_{L_t^{2+\epsilon} L_x^{\frac{6(2+\epsilon)}{\epsilon}}(\Gamma_+(J'))}^{\frac{2+\epsilon}{4}} \|u\|_{L_t^\infty L_x^6(\Gamma_+(J'))}^{1-\frac{2+\epsilon}{4}} \\
&\lesssim E^{\frac{2+\epsilon}{8}} \|u\|_{L_t^\infty L_x^6(\Gamma_+(J'))}^{1-\frac{2+\epsilon}{4}}
\end{aligned}$$

Therefore (7.1) holds.

7.2. Proof of Result 4. By (7.2) we have $E\eta^{-18} \geq 1$. Let n be the largest integer such that $2n \leq 4E\eta^{-18}$. This implies that $n \geq E\eta^{-18}$. Let $A := E\eta^{-18}$. Now we consider the interval $[t_1, A^{2n}t_1] \subset J$. We write $[t_1, A^{2n}t_1] = [t_1, A^2t_1] \cup \dots \cup [A^{2(n-1)}t_1, A^{2n}t_1]$. We have

$$(7.10) \quad \sum_{i=1}^n e(A^{2i}t_1) - e(A^{2(i-1)}t_1) \leq 2E$$

and by the pigeonhole principle there exists $i_0 \in [1, n]$ such that

$$(7.11) \quad e(A^{2i_0}t_1) - e(A^{2(i_0-1)}t_1) \lesssim \eta^{18}$$

Now we choose $a := A^{2(i_0-1)}t_1$ and $b \in [A^{2i_0-1}t_1, A^{2i_0}t_1]$. Let $t'_1 := A^{2(i_0-1)}t_1$, $t'_2 := A^{2i_0-1}t_1$ and $J' := [t'_1, t'_2]$. We apply (3.7) and (7.2) to get

$$\begin{aligned}
(7.12) \quad \|u\|_{L_t^\infty L_x^6(\Gamma_+([t'_1, t'_2]))} &\lesssim \|F(u)\|_{L_t^\infty L_x^1(\Gamma_+([t'_1, t'_2]))} \\
&\lesssim \left(E^{-1}\eta^{18}(E + E^{\frac{1}{3}}) + \eta^{18} + \eta^6 \right)^{\frac{1}{6}} \\
&\lesssim \eta
\end{aligned}$$

8. PROOF OF LEMMA 9

We have after computation of the derivative of $e(t)$

$$(8.1) \quad \partial_t e(t) \geq \int_{|x|=t} F(u) dS$$

and integrating with respect of time

$$(8.2) \quad \int_I \int_{|x| \leq t} g(u) u^6(t', x') dS dt' \lesssim E$$

By using the space and time translation invariance

$$(8.3) \quad \int_J \int_{|x' - x| = |t' - t|} g(u) u^6(t', x') dS dt' \lesssim E$$

Therefore (1.15), (1.22), (8.3) and Hölder inequality

$$\begin{aligned}
(8.4) \quad & \left| - \int_{J'} \frac{\sin(t-t')D}{D} g(u) u^5 dt' \right| \\
&= \left| \frac{1}{4\pi|t-t'|} \int_{|x'-x|=|t'-t|} g^{\frac{5}{6}}(u) u^5 g^{\frac{1}{6}}(u) dS dt' \right| \\
&\lesssim \int_{J'} \frac{1}{|t-t'|} \left(\int_{|x'-x|=|t'-t|} u^6 g(u) dS \right)^{\frac{5}{6}} \left(\int_{|x'-x|=|t'-t|} g(u) dS \right)^{\frac{1}{6}} dt' \\
&\lesssim g^{\frac{1}{6}}(M) \int_{J'} \frac{1}{|t-t'|^{\frac{2}{3}}} \left(\int_{|x'-x|=|t'-t|} u^6 g(u) dS \right)^{\frac{5}{6}} dt' \\
&\lesssim g^{\frac{1}{6}}(M) E^{\frac{5}{6}} \left(\int_{J'} \frac{1}{|t-t'|^4} \right)^{\frac{1}{6}} \\
&\lesssim g^{\frac{1}{6}}(M) \frac{E^{\frac{5}{6}}}{\text{dist}^{\frac{1}{2}}(t, J')}
\end{aligned}$$

Notice that

$$(8.5) \quad u(t) = u_{l, t_i}(t) - \int_{t_i}^t \frac{\sin(t-t')D}{D} u^5(t') g(u(t')) dt'$$

for $i = 1, 2$. We get (3.11) from (8.4) and (8.5).

REFERENCES

- [1] H. Bahouri, P. Gerard, *High frequency approximation of solutions to critical nonlinear wave equations*, Amer. J. Mat 121 (1999), 131-175
- [2] J. Bourgain, *Global well-posedness of defocusing 3D critical NLS in the radial case*, JAMS 12 (1999), 145-171
- [3] J. Ginibre, G. Velo, *Generalized Strichartz inequalities for the wave equation*, Journ. Func. Anal., 133 (1995), 50-68
- [4] M. Grillakis, *Regularity and asymptotic behaviour of the wave equation with a critical nonlinearity*, Ann. of Math. 132 (1990), 485-509
- [5] M. Grillakis, *Regularity for the wave equation with a critical nonlinearity*, Commun. Pure Appl. Math. 45 (1992), 749-774
- [6] L. Kapitanski, *Global and unique weak solutions of nonlinear wave equations*, Math. Res. Letters 1 (1994), 211-223
- [7] M. Keel and T. Tao, *Endpoint Strichartz estimates*, Amer. J. Math, 120 (1998), 955-980
- [8] H. Lindblad, C. D Sogge, *On existence and scattering with minimal regularity for semilinear wave equations*, J. Func. Anal 219 (1995), 227-252
- [9] C. Morawetz, *Time decay for the nonlinear Klein-Gordon equation*, Proc. Roy. Soc. A 306 (1968), 291-296
- [10] J. Rauch, *The u^5 -Klein-Gordon equation*, Nonlinear partial differential equations and their applications (H. Brezis and J. L Lions, eds), Pitman, Boston, 1982, pp 335-364
- [11] J. Shatah, M. Struwe *Geometric Wave Equations*, Courant Lecture Notes in Mathematics 2 (1998)
- [12] J. Shatah, M. Struwe, *Regularity results for nonlinear wave equations*, Annals of Mathematics 138 (1993), 503-518
- [13] J. Shatah, M. Struwe, *Well-Posedness in the energy space for semilinear wave equation with critical growth*, Inter. Math. Research Not., 7 (1994), 303-309
- [14] C. D Sogge, *Lectures on nonlinear wave equations*, Monographs in Analysis II, International Press, 1995
- [15] M. Struwe *Globally regular solutions to the u^5 Klein-Gordon equation*, Ann. Sci. Norm. Sup. Pisa 15 (1988), 495-513
- [16] T. Tao, *Spacetime bounds for the energy-critical nonlinear wave equation in three dimensions*, Dynamics of PDE, 3, 2006, 93-110
- [17] T. Tao, *Global regularity for a logarithmically supercritical defocusing nonlinear wave equation for spherically symmetry data*, J. Hyperbolic Diff. Eq., 4, 2007, 259-266

UNIVERSITY OF CALIFORNIA, LOS ANGELES
E-mail address: `truiroy@math.ucla.edu`